

チャ

カンザワハダニ

発生量は山城で平年比多く、丹波で発生を認めず(平年比やや少ない)、丹後で発生を認めなかった(平年並)。

チャトゲコナジラミ

発生量は山城及び丹波、丹後で例年並。

その他の病害虫

チャノコカクモンハマキ綴葉、幼虫が散見された。

令和6年 チャ害虫調査(カンザワハダニ・50葉調査)

調査日:山城:3/21、24 丹波・丹後:3/24

調査場所	カンザワハダニ							チャトゲコナジラミ					備考	
	寄生 葉率 (%)	100葉当たり			産卵 葉率 (%)	100葉当たり 卵数	発 生 ほ場率 (%)	寄生 葉率 (%)	発生程度別葉数			発生量 指数		発生ほ場 率(%)
		成虫数	幼虫数	合計					0	1	2			
(山城地域)														
宇治市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		調査外でハダニを確認
宇治市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市3	0.0	0	0	0	0.0	0		4.0	48	1	1	3		
宇治市4	2.0	2	0	2	0.0	0		4.0	48	2	0	2		
宇治市5	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治市6	10.0	14	0	14	8.0	298		0.0	50	0	0	0		
城陽市1	6.0	6	0	6	6.0	86		0.0	50	0	0	0		
城陽市2	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
宇治田原町1	0.0	0	0	0	0.0	0		10.0	45	5	0	5		
宇治田原町2	0.0	0	0	0	0.0	0		24.0	38	12	0	12		
宇治田原町3	4.0	4	0	4	2.0	24		0.0	50	0	0	0		
宇治田原町4	0.0	0	0	0	0.0	0		2.0	49	1	0	1		
京田辺市1	2.0	0	12	12	2.0	2		8.0	46	4	0	4		
木津川市1	0.0	0	0	0	0.0	0		6.0	47	3	0	3		
木津川市2	0.0	0	0	0	0.0	0		4.0	48	2	0	2	調査外でハダニを確認	
和束町1	0.0	0	0	0	0.0	0		8.0	46	4	0	4		
和束町2	2.0	4	0	4	2.0	22		34.0	33	15	2	19		
和束町3	2.0	2	0	2	0.0	0		0.0	50	0	0	0		
和束町4	0.0	0	0	0	0.0	0		6.0	47	3	0	3		
和束町5	0.0	0	0	0	0.0	0		4.0	48	2	0	2		
和束町6	2.0	2	0	2	2.0	14		26.0	37	13	0	13		
平均値	1.4	1.6	0.6	2.2	1.0	21.2		6.7	46.7	3.2	0.1	3.5	59.1	
平年比	やや多	やや多	並	やや多	多	多	多	(並)	—	—	—	(並)	(並)	
前年値	0.1	0.1	0.0	0.1	0.0	0.0	4.5	3.3	—	—	—	1.9	40.9	
平年値	0.7	1.1	0.4	1.5	0.2	3.4	19.3	(14.2)				(8.2)	(62.9)	
(丹波地域)														
福知山市1	0.0	0	0	0	0.0	0		4.0	48	2	0	2		チャノコカクモンハマキ綴葉、幼虫散見
福知山市2	0.0	0	0	0	0.0	0		8.0	46	4	0	4		
綾部市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
綾部市2	0.0	0	0	0	0.0	0		2.0	49	1	0	1		
舞鶴市1	0.0	0	0	0	0.0	0		0.0	50	0	0	0		
舞鶴市2	0.0	0	0	0	0.0	0		8.0	46	4	0	4		
平均値	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.7	48.2	1.8	0.0	1.8	66.7	
平年比	やや少	やや少	並	やや少	並	並	やや少	(並)	—	—	—	(並)	(並)	
前年値	0.7	1.3	0.0	1.3	0.3	16.7	33.3	1.3	—	—	—	0.7	33.3	
平年値	0.9	1.1	0.0	1.1	0.1	5.6	25.0	(7.2)				(3.8)	(42.9)	
(丹後地域)														
京丹後市1	0.0	0	0	0	0.0	0		10.0	45	5	0	5		
京丹後市2	0.0	0	0	0	0.0	0		10.0	45	4	1	6		
京丹後市3	0.0	0	0	0	0.0	0		4.0	48	2	0	2		
京丹後市4	0.0	0	0	0	0.0	0		8.0	46	3	1	5		
平均値	0.0	0.0	0.0	0.0	0.0	0.0	0.0	8.0	46.0	3.5	0.5	4.5	100.0	
平年比	並	並	並	並	並	並	並	(並)	—	—	—	(並)	(並)	
前年値	0.0	0.0	0.0	0.0	0.0	0.0	0.0	13.5	—	—	—	6.8	100.0	
平年値	0.7	0.9	0.0	0.9	0.1	1.0	20.0	(19.1)				(10.4)	(78.6)	

カンザワハダニ(3月第5半旬)

地域	平年比	項目 \ 年次	R7	平年比	平年	R6	R5	R4	R3	R2	H31	H30	H29	H28	H27
山城	多	寄生葉率(%)	1.4	やや多	0.7	0.1	0.5	0.9	0.2	0.1	0.1	0.9	0.4	1.2	3.0
		成虫数(100葉当)	1.6	やや多	1.1	0.1	1.0	1.1	0.4	0.1	0.1	0.6	0.4	1.6	5.6
		幼虫数(〃)	0.6	並	0.4	0.0	1.4	0.1	0.0	0.0	0.0	0.4	0.0	0.8	1.0
		合計(〃)	2.2	やや多	1.5	0.1	2.4	1.2	0.4	0.1	0.1	1.0	0.4	2.5	6.6
		産卵葉率(%)	1.0	多	0.2	0.0	0.4	0.6	0.1	0.1	0.0	0.1	0.0	0.5	0.4
		卵数(100葉当)	21.2	多	3.4	0.0	10.8	8.9	4.4	0.4	0.0	0.9	0.0	8.0	1.1
		発生ほ場率(%)	45.5	多	19.3	4.5	22.7	22.7	9.1	4.8	4.5	27.3	13.6	36.4	47.6
丹波	やや少	寄生葉率(%)	0.0	やや少	0.9	0.7	1.3	0.0	0.3	1.0	0.3	0.3	0.0	1.0	4.3
		成虫数(100葉当)	0.0	やや少	1.1	1.3	1.3	0.0	0.3	1.3	1.3	0.3	0.0	1.0	4.3
		幼虫数(〃)	0.0	並	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		合計(〃)	0.0	やや少	1.1	1.3	1.3	0.0	0.3	1.3	1.3	0.3	0.0	1.0	4.3
		産卵葉率(%)	0.0	並	0.1	0.3	0.0	0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.3
		卵数(100葉当)	0.0	並	5.6	16.7	0.0	0.0	0.0	0.0	33.3	5.3	0.0	0.0	0.7
		発生ほ場率(%)	0.0	やや少	25.0	33.3	50.0	0.0	16.7	33.3	16.7	16.7	0.0	33.3	50.0
丹後	並	寄生葉率(%)	0.0	並	0.7	0.0	1.0	0.0	0.5	1.5	0.0	0.0	0.0	0.5	3.0
		成虫数(100葉当)	0.0	並	0.9	0.0	2.5	0.0	0.5	2.0	0.0	0.0	0.0	0.5	3.5
		幼虫数(〃)	0.0	並	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
		合計(〃)	0.0	並	0.9	0.0	2.5	0.0	0.5	2.0	0.0	0.0	0.0	0.5	3.5
		産卵葉率(%)	0.0	並	0.1	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0
		卵数(100葉当)	0.0	並	1.0	0.0	0.0	0.0	10.0	0.0	0.0	0.0	0.0	0.0	0.0
		発生ほ場率(%)	0.0	並	20.0	0.0	50.0	0.0	25.0	75.0	0.0	0.0	0.0	25.0	25.0

チャトゲコナジラミ(3月第5半旬)

地域	平年(例年)比	項目 \ 年次	R7	平年(例年)比	平年(例年)	R6	R5	R4	R3	R2	H31	H30	H29	H28	H27
山城	(並)	寄生葉率(%)	6.7	(並)	(14.2)	3.3	4.7	7.0	14.2	34.3	19.5	16.73	—	—	—
		発生量指数	3.5	(並)	(8.2)	1.9	2.6	3.7	7.5	23.7	10.0	7.66	—	—	—
		発生ほ場率(%)	59.1	(並)	(62.9)	40.9	54.5	50.0	63.6	85.7	77.3	68.20	—	—	—
丹波	(並)	寄生葉率(%)	3.7	(並)	(7.2)	1.3	0.0	0.0	2.0	22.0	21.3	4.00	—	—	—
		発生量指数	1.8	(並)	(3.8)	0.7	0.0	0.0	1.0	12.0	11.0	2.00	—	—	—
		発生ほ場率(%)	66.7	(並)	(42.9)	33.3	0.0	0.0	16.7	100.0	100.0	50.00	—	—	—
丹後	(並)	寄生葉率(%)	8.0	(並)	(19.1)	13.5	6.5	3.0	12.5	47.0	44.5	6.50	—	—	—
		発生量指数	4.5	(並)	(10.4)	6.8	3.8	1.5	6.5	28.8	22.5	3.25	—	—	—
		発生ほ場率(%)	100.0	(並)	(78.6)	100.0	75.0	25.0	75.0	100.0	100.0	75.00	—	—	—